arXiv:1304.1872v1 [math.NT] 6 Apr 2013

HOMOGENEOUS SPACES OF HILBERT TYPE

MIKHAIL BOROVOI

ABSTRACT. Let k be a global field. Let G be a connected linear algebraic k-group, assumed reductive when k is a function field. It follows from a result of the preprint [BFP] by Bary-Soroker, Fehm and Petersen that when H is a smooth connected k-subgroup of G, the quotient space G/H is of Hilbert type. We prove a similar result for certain non-connected k-subgroups H of G. In particular, we prove that if G is a simply connected k-group over a number field k, and H is an abelian k-subgroup of G, not necessarily connected, then G/H is of Hilbert type.

1. INTRODUCTION

Let k be a field, and let X be a geometrically integral k-variety. For the definition of a *thin subset* $\Theta \subset X(k)$ we refer to Serre [Se, Def. 3.1.1], see also Bary-Soroker, Fehm and Petersen [BFP, Section 2]. If $\Theta' \subset \Theta \subset X(k)$ and Θ is thin, then Θ' is thin too. If we have a finite family of thin subsets $\Theta_1, \dots, \Theta_r \subset X(k)$, then their union $\Theta_1 \cup \dots \cup \Theta_r$ is again a thin subset. The notion of a variety *of Hilbert type* was introduced by Colliot-Thélène and Sansuc [CS]. By definition, see Serre [Se, Def. 3.1.2], a k-variety X is of Hilbert type if $X(k)$ is not thin. A field k is called Hilbertian if the affine line $\mathbf{A}^1_k$ is of Hilbert type. All the global fields (number fields and function fields) are Hilbertian, see [Se, Thm. 3.4.1] and Fried and Jarden [FJ, Thm. 13.4.2].

It was proved by Colliot-Thélène and Sansuc in [CS, Cor. 7.15] that any connected reductive algebraic k-group over a Hilbertian field k is of Hilbert type. Moreover, we have the following more general result, cf. [BFP]:

Proposition 1.1. *Let k be a Hilbertian field. Let G be a smooth connected linear algebraic k-group, assumed reductive when k is not perfect. Let $H \subset G$ be a smooth connected k-subgroup. Then the quotient space G/H is a k-variety of Hilbert type.*

Proof. When k is perfect, see Bary-Soroker, Fehm and Petersen [BFP, Cor. 4.5]. When k is not perfect (and G is reductive), the proof is similar. ☐

In this note we consider the case when k is a *global field* and G is a smooth connected linear algebraic k-group. We prove that G/H is of Hilbert type for certain *non-connected and non-smooth* subgroups $H \subset G$. In particular, we prove the following theorem:

Theorem 1.2. *Let G be a simply connected k-group over a number field k, and let $H \subset G$ be an abelian k-subgroup, not necessarily connected. Then G/H is of Hilbert type.*

Proof. This theorem is a special case of Theorem 2.2 below. ☐

2010 *Mathematics Subject Classification.* Primary: 14M17, 12E25, Secondary: 20G30, 12E30.

Key words and phrases. Hilbertian field, variety of Hilbert type, linear algebraic group, homogeneous space.

Partially supported by the Hermann Minkowski Center for Geometry.

The plan of the rest of the note is as follows. In the end of the Introduction we give some notation. In Section 2 we consider homogeneous spaces over a number field. The main result of this section, Theorem 2.2, implies Theorem 1.2, and its proof uses [BFP, Thm. 1.1]. In Section 3 we consider homogeneous spaces over a global function field. The main result of this section is Theorem 3.1, the proof of which is similar to and easier than that of Theorem 2.2, and uses [BFP, Cor. 1.2]. In Section 4 we give an alternative proof of Theorem 2.2. We show that a homogeneous space G/H as in Theorem 2.2 has the WWA (weak weak approximation) property, cf. [Se, Def. 3.5.6], and therefore, by a theorem of Ekedahl and Colliot-Thélène (see [Se, Thm. 3.5.7]) G/H is of Hilbert type.

Notation 1.3. Let k be a field. By $\bar{k}$ we denote a fixed separable closure of k. If X is a k-variety, we write $\overline{X} := X \times_k \bar{k}$. If H is a linear algebraic group (not necessarily connected or smooth), we write $H^{\rm mult}$ for the largest quotient group of H which is a group of multiplicative type.

Let G be a smooth connected linear algebraic k-group, and $H \subset G$ a k-subgroup, not necessarily connected or smooth. We write G/H for the quotient variety. For the existence of G/H see [Sp, Thm. 12.2.1] in the case when H is smooth, and [DG, Exp. VI$_A$, Thm. 3.2(i)] in the general case. Since G is smooth, so is G/H, see [CGP, after Example A.1.12, bottom of p. 395].

If k is a number field, we denote by $\mathscr{V}(k)$ the set of all places of k, and by $\mathscr{V}_\infty(k)$ the set of all archimedean places. If $v \in \mathscr{V}(k)$, we write k_v for the completion of k at v. We set $k_\infty := \prod_{v\in\mathscr{V}_\infty(k)} k_v$. If X is a k-variety, then $X(k)$ embeds into $X(k_\infty) = \prod_{v\in\mathscr{V}_\infty(k)} X(k_v)$.

2. NUMBER FIELDS

Notation 2.1. Let k be a field of characteristic 0. Let H be a linear k-group. We assume that the kernel $H_1 := \ker[H \to H^{\rm mult}]$ is connected and geometrically character-free. In this case H_1 is an extension of a connected semisimple group by a unipotent group. Note that in characteristic 0, any *connected* linear k-group H and any *abelian* linear k-group H satisfy this assumption.

Let G be a connected linear algebraic k-group. We use the following notation:
$G^{\rm u}$ is the unipotent radical of G;
$G^{\rm red} = G/G^{\rm u}$, it is reductive;
$G^{\rm ss} = [G^{\rm red}, G^{\rm red}]$, it is semisimple;
$G^{\rm tor} = G^{\rm red}/G^{\rm ss}$, it is a torus;
$G^{\rm ssu} = \ker[G \to G^{\rm tor}]$, it is an extension of the semisimple group $G^{\rm ss}$ by the unipotent group $G^{\rm u}$.

Note that $G^{\rm tor}$ is the largest quotient torus of G and that $G^{\rm mult} = G^{\rm tor}$.

Theorem 2.2. *Let k be a number field. Let G be a connected linear algebraic k-group, and let $H \subset G$ be a k-subgroup, not necessarily connected. We assume that $G^{\rm ss}$ is simply connected and that $H_1 := \ker[H \to H^{\rm mult}]$ is connected and geometrically character-free. Then the variety $X := G/H$ is of Hilbert type.*

We need two lemmas.

Lemma 2.3. *Let F be a linear algebraic group over a field k, and let Γ be a subgroup of finite index of $F(k)$. If F is of Hilbert type, then the set $\Gamma \subset F(k)$ is not thin.*

Proof. Since Γ is of finite index in $F(k)$, we have $F(k) = f_1\Gamma \cup \cdots \cup f_m\Gamma$ for some finite set $f_1, \ldots, f_m \subset F(k)$. Suppose for the sake of contradiction that Γ is thin, then all the sets

$f_i\Gamma$ are thin $(i=1,\dots,m)$, hence the finite union $F(k)=f_1\Gamma\cup\dots\cup f_m\Gamma$ is thin, which contradicts to the assumption that F is of Hilbert type. □

Lemma 2.4. *Let T be a k-torus over a number field k. Let $T(k_\infty)^\circ$ denote the identity component of $T(k_\infty)$. Then the set $T(k)\cap T(k_\infty)^\circ$ is not thin.*

Proof. Since the number field k is Hilbertian, the torus T is a variety of Hilbert type, cf. [CS, Cor. 7.14]. Consider the group of connected components $\pi_0(T(k_\infty)):=T(k_\infty)/T(k_\infty)^\circ$, it is finite. It follows that the subgroup $T(k)\cap T(k_\infty)^\circ$ is of finite index in $T(k)$. Since T is of Hilbert type, by Lemma 2.3 $T(k)\cap T(k_\infty)^\circ$ is not thin. □

Proof of Theorem 2.2. We prove the theorem in two steps, using the method of [B1].

Step 1. The inclusion $H\hookrightarrow G$ induces a homomorphism $H^{\mathrm{mult}}\to G^{\mathrm{tor}}$, which need not be injective. Choose an embedding $j\colon H^{\mathrm{mult}}\hookrightarrow Q$ into a quasi-trivial k-torus Q, and denote by $m\colon H\to H^{\mathrm{mult}}$ the canonical epimorphism. Set $G_Y:=G\times_k Q$, then we have a diagonal embedding

$$H\hookrightarrow G_Y,\quad h\mapsto (h,j(m(h)))\in G\times_k Q=G_Y.$$

Let H_Y denote the image of H in G_Y, then $(H_Y)^{\mathrm{mult}}$ embeds into $(G_Y)^{\mathrm{tor}}$. Set $Y=G_Y/H_Y$. We have a map

$$\pi\colon Y\to X,\ (g,q)H_Y\mapsto gH,\text{where } (g,q)\in G\times_k Q=G_Y.$$

The subgroup $Q\subset G_Y$ acts on Y on the left, and Y is a left torsor over X under Q. It follows that all the fibers of π are geometrically integral, in particular, the generic fiber of π is geometrically integral. Therefore, by [CS, Prop. 7.13], in order to prove that X is of Hilbert type, it suffices to show that Y is of Hilbert type.

Step 2. From now on we assume that H^{mult} embeds into G^{tor}, and we regard H^{mult} as a subgroup of G^{tor}. The canonical homomorphism $G\to G^{\mathrm{tor}}$ takes H to $H^{\mathrm{mult}}\subset G^{\mathrm{tor}}$ and induces a G-equivariant map $\mu\colon X=G/H\to T:=G^{\mathrm{tor}}/H^{\mathrm{mult}}$. We see that T is a k-torus. We have a commutative diagram

$$\begin{tikzcd}
G \arrow[r] \arrow[d, "t"'] \arrow[dr, "\phi"] & X \arrow[d, "\mu"] \\
G^{\mathrm{tor}} \arrow[r] & T
\end{tikzcd}$$

where the horizontal arrows are the canonical maps $G\to X:=G/H$ and $G^{\mathrm{tor}}\to T:=G^{\mathrm{tor}}/H^{\mathrm{mult}}$. Note that the diagonal arrow $\phi\colon G\to T$ is a homomorphism of k-groups. We see from the diagram $\mu(X(k_\infty))\supset\phi(G(k_\infty))$. On the other hand, the homomorphism ϕ is surjective, hence it is a smooth map, and therefore the subgroup $\phi(G(k_\infty))\subset T(k_\infty)$ is open, hence it contains $T(k_\infty)^\circ$. Thus $\mu(X(k_\infty))$ contains $T(k_\infty)^\circ$.

Let $t\in T(k)\cap T(k_\infty)^\circ$ be a k-point. We show that the fiber $X_t:=\mu^{-1}(t)$ is of Hilbert type. First we prove that X_t has a k-point. Clearly X_t is a homogeneous space of $G^{\mathrm{ssu}}:=\ker[G\to G^{\mathrm{tor}}]$. From the assumption that G^{ss} is simply connected we deduce that G^{ssu} is simply connected. Let $x_0\in X(k)$ denote the image of the unit element of G in $X(k)=G(k)/H(k)$, then the stabilizer of x_0 in G^{ssu} is $H\cap G^{\mathrm{ssu}}=H_1$, because H^{mult} embeds into G^{tor}. Let $\overline{x}\in X_t(\overline{k})$ and let $\overline{H}_{1,\overline{x}}$ denote the stabilizer of $\overline{x}$ in $\overline{G^{\mathrm{ssu}}}$, then $\overline{H}_{1,\overline{x}}$ is conjugate to $\overline{H_1}$ in $\overline{G}$, hence it is connected and character-free. Since $t\in T(k)\cap T(k_\infty)^\circ\subset\mu(X(k_\infty))$, we see that the homogeneous space X_t has a k_v-point for every $v\in\mathcal{V}_\infty(k)$. By a local-global principle for such homogeneous spaces, see [B1, Prop. 3.4(i)], the homogeneous space X_t has a k-point x. Now let $H_{1,x}$ denote the stabilizer of x in G^{ssu}, then $H_{1,x}$ is connected. Since X_t is isomorphic to $G^{\mathrm{ssu}}/H_{1,x}$, where $H_{1,x}$ is a connected k-subgroup of a connected linear k-group G^{ssu}, by Proposition 1.1 X_t is of Hilbert type.

By Lemma 2.4 the subset $T(k)\cap T(k_\infty)^\circ \subset T(k)$ is not thin, and we have shown that for any $t\in T(k)\cap T(k_\infty)^\circ$ the fiber X_t of X over t is of Hilbert type. By [BFP, Theorem 1.1], the variety X is of Hilbert type. ☐

3. FUNCTION FIELDS

Let k be a field of characteristic $p>0$. Let G be a smooth connected *reductive* linear algebraic group over k. We set $G^{\mathrm{ss}}=[G,G]$ and $G^{\mathrm{tor}}=G/G^{\mathrm{ss}}$.

Recall that a global function field is the function field of a smooth, projective, connected curve over a finite field.

Theorem 3.1. *Let k be a global function field. Let G be a smooth, connected, reductive k-group, and let $H\subset G$ be a k-subgroup, not necessarily connected or smooth. We assume that G^{ss} is simply connected and that $H_1:=\ker[H\to H^{\mathrm{mult}}]$ is smooth, connected and semisimple. Then the quotient variety $X:=G/H$ is of Hilbert type.*

Proof. *Step 1.* As in the proof of Theorem 2.2, we reduce the theorem to the case when H^{mult} embeds into G^{tor}.

Step 2. From now on we assume that H^{mult} embeds into G^{tor}, and we regard H^{mult} as a subgroup of G^{tor}. The canonical homomorphism $G\to G^{\mathrm{tor}}$ takes H to $H^{\mathrm{mult}}\subset G^{\mathrm{tor}}$ and induces a G-equivariant map $\mu\colon X\to T:=G^{\mathrm{tor}}/H^{\mathrm{mult}}$. We see that T is a k-torus, hence T is of Hilbert type, cf. [CS, Cor. 7.14].

Let $t\in T(k)$ be a k-point. We show that the fiber $X_t:=\mu^{-1}(t)$ is of Hilbert type. Clearly X_t is a homogeneous space of $G^{\mathrm{ss}}:=\ker[G\to G^{\mathrm{tor}}]$. Let $\overline{x}\in X_t(\overline{k})$. Since H^{mult} embeds into G^{tor}, the stabilizer $\overline{H}_{1,\overline{x}}$ of $\overline{x}$ in $\overline{G^{\mathrm{ss}}}$ is conjugate in $\overline{G}$ to $\overline{H}\cap\overline{G^{\mathrm{ss}}}=\overline{H_1}$, hence it is smooth, connected and semisimple. Now by Proposition 3.2 below, the homogeneous space X_t has a k-point x. Let $H_{1,x}$ denote the stabilizer of x in G^{ss}, then $H_{1,x}$ is smooth and connected. Since X_t is isomorphic to $G^{\mathrm{ss}}/H_{1,x}$ where $H_{1,x}$ is a smooth connected k-subgroup of a smooth connected reductive k-group G^{ss}, by Proposition 1.1 X_t is of Hilbert type.

Since T is of Hilbert type and for any $t\in T(k)$ the fiber X_t of X over t is of Hilbert type, by [BFP, Cor. 1.2] the variety X is of Hilbert type. ☐

In the proof of Theorem 3.1 we used the following result for function fields, which is similar to [B1, Prop. 3.4(i)] for number fields.

Proposition 3.2. *Let k be a global function field. Let G be a smooth, connected, semisimple, simply connected k-group. Let X be a right homogeneous space of G. Let $\overline{H}$ be the stabilizer of a geometric point $\overline{x}\in X(\overline{k})$, where $\overline{k}$ is a fixed separable closure of k. Assume that $\overline{H}$ is smooth, connected and semisimple. Then X has a k-point.*

Proof. For a commutative étale algebra L/K, we consider the category of pairs $(P_{(L)},\alpha_{(L)})$, where $P_{(L)}$ is a right torsor over L of $G_L:=G\times_k L$ and $\alpha_{(L)}\colon P_{(L)}\to X_L$ is a G_L-equivariant L-morphism. This category is clearly a groupoid (maybe empty). Over the separable closure $\overline{k}$ of k such a pair clearly exists and any two such pairs are isomorphic. We obtain a fibered category $\mathscr{G}_X$ over $\operatorname{Spec} k$, which is a gerbe locally bound by $\overline{H}$, see Giraud [G, Ch. IV, 5.1.2]. Since $\overline{H}$ is semisimple, the band of $\mathcal{G}$ can be represented be a quasi-split k-form H_0 of $\overline{H}$, see [D1, Lemma 1.1], and the gerbe $\mathcal{G}$ is locally bound by H_0. By definition, the gerbe $\mathcal{G}_X$ is neutral (trivial) if a pair (P,α) as above exists over k, see Giraud [G, Ch. IV, Prop. 5.1.4(ii)]. For details see also [D3, § 1]. Since k is a global function field, and since H_0 is smooth, connected and semisimple, by Douai's theorem [D2, Cor. 5.2] any gerbe over k which is locally bound by H_0 is neutral. Thus our gerbe $\mathcal{G}_X$ is neutral, i.e., there exists pair (P,α) as above, defined over k. By Harder's theorem [H, Satz A] any torsor of a simply connected semisimple k-group over a global function field k has a k-point. Thus P has a k-point p. We obtain a k-point of X by taking $x=\alpha(p)\in X(k)$. This completes the proofs of Proposition 3.2 and Theorem 3.1. ☐

4. WEAK WEAK APPROXIMATION

In this section we give an alternative proof of Theorem 2.2, using the WWA (weak weak approximation) property. In this section k is always a number field and $S\subset\mathcal{V}(k)$ is a finite subset.

One says that a k-variety X has the WWA (weak weak approximation) property, if there exists a finite subset $S_0\subset\mathcal{V}(k)$ with the following property: for any finite subset $S\subset\mathcal{V}(k)$ such that $S\cap S_0=\emptyset$, the variety X has weak approximation in S (i.e. $X(k)$ is dense in $\prod_{v\in S}X(k_v)$). By a theorem of T. Ekedahl and J.-L. Colliot-Thélène, see [E] and [Se, Thm. 3.5.7], any variety with the WWA property is of Hilbert type. We shall prove that G/H as in Theorem 2.2 has *WWA*, thus it is of Hilbert type.

Notation 4.1. Let X be a smooth k-variety. We write $\mathrm{Br}(X)$ for the cohomological Brauer group of X, i.e. $\mathrm{Br}(X)=H^2_{\text{ét}}(X,\mathbf{G}_\mathrm{m})$. We set $\mathrm{Br}_1(X)=\ker[\mathrm{Br}(X)\to\mathrm{Br}(\overline{X})]$ where $\overline{X}=X\times_k\overline{k}$. We define the *algebraic Brauer group* $\mathrm{Br}_\mathrm{a}(X)$ by $\mathrm{Br}_\mathrm{a}(X)=\mathrm{coker}[\mathrm{Br}(k)\to\mathrm{Br}_1(X)]$. For a finite subset $S\subset\mathcal{V}(k)$ we set

$$Б_S(X)=\ker\left[\mathrm{Br}_\mathrm{a}(X)\to\prod_{v\notin S}\mathrm{Br}_\mathrm{a}(X_{k_v})\right].$$

Set $Б_{S,\emptyset}(X)=Б_S(X)/Б_\emptyset(X)$.

Let B be a discrete $\mathrm{Gal}(\overline{k}/k)$-module which is finitely generated as an abelian group (we say just "a finitely generated Galois module"). We write

$$Ш^i_S(k,B)=\ker\left[H^i(k,B)\to\prod_{v\notin S}H^i(k_v,B)\right],$$

where $i=1,2,\dots$. We set $Ш^i_{S,\emptyset}(k,B):=Ш^i_S(k,B)/Ш^i_\emptyset(k,B)$.

Let $A\to B$ be a morphism of finitely generated $\mathrm{Gal}(\overline{k}/k)$-modules. We write $\mathbb{H}^i(k,A\to B)$ $(i=1,2,\dots)$ for i-th Galois hypercohomology of the complex $A\to B$, where A is in degree 0 and B is in degree 1. We define $Ш^i_S(k,A\to B)$ and $Ш^i_{S,\emptyset}(k,A\to B)$ as above.

Lemma 4.2. *Let k be a number field, and let $A\to B$ be a homomorphism of finitely generated $\mathrm{Gal}(\overline{k}/k)$-modules, where A is free as an abelian group. Let K/k be the finite Galois extension in $\overline{k}$ corresponding to $\ker\left[\mathrm{Gal}(\overline{k}/k)\to\mathrm{Aut}\,A\times\mathrm{Aut}\,B\right]$. Let $S\subset\mathcal{V}(k)$ be a finite subset formed by places with cyclic decomposition groups in $\mathrm{Gal}(K/k)$. Then $Ш^2_{S,\emptyset}(k,A\to B)=0$.*

Proof. Set $\Gamma = \mathrm{Gal}(K/k)$. Choose an epimorphism of Γ-modules $P \twoheadrightarrow B$, where P is a permutation Γ-module. Set $L = A \times_B P$, then we have a quasi-isomorphism $(L \to P) \to (A \to B)$, and therefore isomorphisms $\mathbb{H}^2(k, L \to P) \xrightarrow{\sim} \mathbb{H}^2(k, A \to B)$ and $\text{Ш}^2_{S,\emptyset}(k, L \to P) \xrightarrow{\sim} \text{Ш}^2_{S,\emptyset}(k, A \to B)$.

We have a short exact sequence of complexes

$$0 \to (0 \to P) \to (L \to P) \to (L \to 0) \to 0,$$

which induces a hypercohomology exact sequence

$$0 = H^1(k, P) \to \mathbb{H}^2(k, L \to P) \to H^2(k, L) \to H^2(k, P),$$

which in turn induces isomorphisms

$$\text{Ш}^2_S(k, L \to P) \xrightarrow{\sim} \text{Ш}^2_S(k, L) \quad \text{and} \quad \text{Ш}^2_{S,\emptyset}(k, L \to P) \xrightarrow{\sim} \text{Ш}^2_{S,\emptyset}(k, L)$$

(because $\text{Ш}^2_S(k, P) = 0$ by [Sa, (1.9.1)]).

Note that L is a free abelian group. We set $L^\vee := \mathrm{Hom}(L, \mathbb{Z})$. Choose an epimorphism $P' \twoheadrightarrow L^\vee$, where P' is a permutation Γ-module. We obtain an embedding $L \hookrightarrow P''$, where $P'' := (P')^\vee$ is a permutation Γ-module. Set $C = P''/L$. The short exact sequence of Γ-modules

$$0 \to L \to P'' \to C \to 0$$

induces a cohomology exact sequence

$$0 = H^1(k, P'') \to H^1(k, C) \to H^2(k, L) \to H^2(k, P''),$$

which in turn induces isomorphisms

$$\text{Ш}^1_S(k, C) \xrightarrow{\sim} \text{Ш}^2_S(k, L) \quad \text{and} \quad \text{Ш}^1_{S,\emptyset}(k, C) \xrightarrow{\sim} \text{Ш}^2_{S,\emptyset}(k, L)$$

(because $\text{Ш}^2_S(k, P'') = 0$ by [Sa, (1.9.1)]).

Since $\mathrm{Gal}(\bar{k}/K)$ acts trivially on C, and S consists of places with cyclic decomposition groups in $\mathrm{Gal}(K/k)$, by [B2, Cor. 3.3] $\text{Ш}^1_{S,\emptyset}(k, C) = 0$. Thus $\text{Ш}^2_{S,\emptyset}(k, A \to B) = 0$, which proves the lemma. $\square$

Alternative proof of Theorem 2.2. Let G and H be as in Theorem 2.2. We have a homomorphism of $\mathrm{Gal}(\bar{k}/k)$-modules $\widehat{G} \to \widehat{H}$, where $\widehat{G}$ and $\widehat{H}$ are the corresponding geometric character groups. Let K/k be the finite Galois extension in $\bar{k}$ corresponding to the kernel

$$\ker\left[\mathrm{Gal}(\bar{k}/k) \to \mathrm{Aut}\,\widehat{G} \times \mathrm{Aut}\,\widehat{H}\right].$$

Let $S_0 \subset \mathcal{V}(k)$ denote the (finite) set of all places of k with *non-cyclic* decomposition groups in $\mathrm{Gal}(K/k)$. Let $S \subset \mathcal{V}(k)$ be a finite set such that $S \cap S_0 = \emptyset$.

By [BvH, Thm. 7.2] there is a canonical isomorphism $\mathrm{Br}_{\mathrm{a}}(G/H) \xrightarrow{\sim} \mathbb{H}^2(k, \widehat{G} \to \widehat{H})$, which induces a canonical isomorphism $\text{Б}_{S,\emptyset}(G/H) \xrightarrow{\sim} \text{Ш}^2_{S,\emptyset}(k, \widehat{G} \to \widehat{H})$. By Lemma 4.2 $\text{Ш}^2_{S,\emptyset}(k, \widehat{G} \to \widehat{H}) = 0$, hence $\text{Б}_{S,\emptyset}(G/H) = 0$.

The Brauer-Manin obstruction of [B1] to weak approximation in S for a k-variety X is a certain map

$$m_S \colon \prod_{v \in S} X(k_v) \to \text{Б}_{S,\emptyset}(G/H)^D,$$

where $\text{Б}_{S,\emptyset}(G/H)^D := \mathrm{Hom}(\text{Б}_{S,\emptyset}(G/H), \mathbb{Q}/\mathbb{Z})$. Since $\text{Б}_{S,\emptyset}(G/H) = 0$, we obtain that $\text{Б}_{S,\emptyset}(G/H)^D = 0$, hence m_S is identically zero. We see that the group G^{ss} is simply connected, $H_1 := \ker[H \to H^{\mathrm{mult}}]$ is connected and geometrically character-free, and the obstruction m_S is identically zero, hence by [B1, Thm. 2.3] the variety G/H has weak approximation in S.

We have proved that G/H has WWA, hence it is of Hilbert type. $\square$

Acknowledgements. The author is grateful to Lior Bary-Soroker, Jean-Claude Douai, and Arno Fehm for very helpful discussions.

REFERENCES

[BFP] L. Bary-Soroker, A. Fehm and S. Petersen, *On varieties of Hilbert type,* `arXiv:1302.4038v2[math.AG]`.

[B1] M. Borovoi, *The Brauer-Manin obstructions for homogeneous spaces with connected or abelian stabilizer,* J. reine angew. Math. **473** (1996), 181–194.

[B2] M. Borovoi, *Vanishing of algebraic Brauer-Manin obstructions,* J. Ramanujan Math. Soc. **26** (2011), 333–349.

[BvH] M. Borovoi and J. van Hamel, *Extended equivariant Picard complexes and homogeneous spaces,* Transform. Groups **17** (2012), 51–86.

[CS] J.-L. Colliot-Thélène and J.-J. Sansuc, *Principal homogeneous spaces under flasque tori: applications,* J. Algebra **106** (1987), 148–205.

[CGP] B. Conrad, O. Gabber, G. Prasad, *Pseudo-reductive groups,* New Mathematical Monographs, 17, Cambridge University Press, Cambridge, 2010.

[DG] M. Demazure et A. Grothendieck (Eds.): *Schémas en groupes.* Séminaire de Géométrie Algébrique du Bois Marie 1962-64 (SGA 3). Augmented and corrected re-edition of the original by P. Gille and P. Polo. Tome 1, Documents mathématiques 7, Société Mathématique de France, 2011.

[D1] J.-C. Douai, *2-cohomologie galoisienne des groupes semi-simples définis sur les corps locaux,* C. R. Acad. Sci. Paris Sér. A-B **280** (1975), A321–A323.

[D2] J.-C. Douai, *Cohomologie galoisienne des groupes semi-simples définis sur les corps globaux,* C. R. Acad. Sci. Paris Sér. A-B **281** (1975), A1077–A1080.

[D3] J.-C. Douai, *Espaces homogènes et arithmétique des schémas en groupes réductifs sur les anneaux de Dedekind,* J. Théor. Nombres Bordeaux **7** (1995), 21–26.

[E] T. Ekedahl, *An effective version of Hilbert's irreducibility theorem,* Séminaire de Théorie des Nombres, Paris 1988–1989, in: Progr. Math. 91, Birkhäuser, Boston, MA, 1990, pp. 241–249.

[FJ] M. D. Fried and M. Jarden, *Field arithmetic,* 3rd ed., Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, 11, Springer-Verlag, Berlin, 2008.

[G] J. Giraud, *Cohomologie non abélienne,* Die Grundlehren der mathematischen Wissenschaften, Band 179, Springer-Verlag, Berlin-New York, 1971.

[H] G. Harder, *Über die Galoiskohomologie halbeinfacher algebraischer Gruppen, III,* J. reine angew. Math. **274/275** (1975), 125–138.

[Sa] J.-J. Sansuc, *Groupe de Brauer et arithmétique des groupes algébriques linéaires sur un corps de nombres*, J. reine angew. Math. **327** (1981), 12–80.

[Se] J.-P. Serre, *Topics in Galois theory,* Research Notes in Mathematics 1, Jones and Bartlett Publishers, Boston, MA, 1992.

[Sp] T. A. Springer, *Linear algebraic groups,* 2nd edition, Progress in Mathematics, 9, Birkhäuser, Boston, MA, 1998.

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, 69978 TEL AVIV, ISRAEL

E-mail address: `borovoi@post.tau.ac.il`